

6420 Rockledge Drive, Suite 4200, Bethesda, MD 20817 P: 240.487.7522 / F: 360.597.1464

Medications to Avoid Prior to Surgery

- **DO NOT TAKE** any medications that contain **Aspirin** or any of the medications you take asneeded that are listed below for **5 DAYS** before your surgery.
- If you take daily blood-thinning medications such as Plavix, Xarelto, Coumadin, or any others, you must also stop these at least <u>5 DAYS</u> prior to surgery. You should do this **only** after discussing with your cardiologist or primary care physician a plan on stopping and restarting these medications.
- Avoiding these medications is very important in preventing bleeding complications during and after surgery
- If you forget to stop these medications, please contact us at the above number **IMMEDIATELY**.
- Taking Tylenol before surgery is okay.

DO NOT TAKE:

Aspirin (ASA) Coumadin Naprosyn (naproxen)

AdvilEcotrinNuprinAggrenoxEliquisPamprinAleveExcedrinPradaxa

Alka Seltzer Effervescent Fioricet Relafen (nabumetone)
Anacin Ibuprofen Sinutab/Sine-Aid
Anaprox (naproxen) Indocin or Maxalt Soma Compound

Baby Aspirin Midol Sulindac Bayer Aspirin Mobic (meloxicam) Toradol

Brilinta Motrin Voltaren (diclofenac)

Bufferin Non-steroidal anti- Warfarin Carisoprodol Compound inflammatories Xarelto

Herbal Supplements to avoid: Echinacea, Ephedra, Garlic, Ginger, Ginko, Ginko Biloba, Ginseng, Kava, St. John's Wort